

**GCE**

**Chemistry A**

Unit **H032/01**: Breadth in chemistry

Advanced Subsidiary GCE

**Mark Scheme for June 2016**

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.















All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2016

Annotations available in RM Assessor

Annotation	Meaning
	Correct response
	Incorrect response
	Omission mark
	Benefit of doubt given
	Contradiction
	Rounding error
	Error in number of significant figures
	Error carried forward
	Level 1
	Level 2
	Level 3
	Benefit of doubt not given
	Noted but no credit given
	Ignore

Abbreviations, annotations and conventions used in the detailed Mark Scheme (to include abbreviations and subject-specific conventions).

<b>Annotation</b>	<b>Meaning</b>
/, OR	alternative and acceptable answers for the same marking point
✓	Separates marking points
<b>DO NOT ALLOW</b>	Answers which are not worthy of credit
<b>IGNORE</b>	Statements which are irrelevant
<b>ALLOW</b>	Answers that can be accepted
( )	Words which are not essential to gain credit
—	Underlined words must be present in answer to score a mark
<b>ECF</b>	Error carried forward
<b>AW</b>	Alternative wording
<b>ORA</b>	Or reverse argument

## SECTION A

Question	Answer	Marks	AO element	Guidance
1	B	1		
2	C	1		
3	D	1		
4	A	1		
5	C	1		<b>ALLOW +5 OR 5+ in box</b>
6	C	1		<b>ALLOW 8 in box</b>
7	B	1		
8	D	1		
9	C	1		
10	B	1		
11	D	1		
12	B	1		
13	B	1		
14	C	1		
15	B	1		
16	D	1		
17	D	1		
18	B	1		
19	B	1		
20	D	1		
	<b>Total</b>	<b>20</b>		

## SECTION B

Question			Answer	Marks	AO element	Guidance
21	(a)	(i)	<p><b>Similarities:</b> (Same) <b>number</b> of protons <b>AND</b> electrons ✓</p> <p><b>Differences:</b> (Different) <b>number</b> of neutrons ✓</p>	2	AO1.1 x2	<p><b>ALLOW</b> same electron configuration</p> <p><b>ALLOW</b> 'amount' for 'number'</p> <p><b>IGNORE</b> different masses/mass numbers (<i>Question asks for atomic structures</i>)</p>
	(b)	(i)	<p><b>FIRST CHECK ANSWER ON THE ANSWER LINE</b> <b>If answer = 63.62 award 2 marks</b></p> <p>-----</p> <p><math display="block">\frac{(63 \times 69.17) + (65 \times 30.83)}{100}</math></p> <p><b>OR</b> 63.6166 <b>OR</b> 63.617 ✓</p> <p>= 63.62 (to 2 DP) ✓</p> <p><b>IGNORE</b> any units with <math>A_r</math></p>	2	AO1.2 x2	<p><b>ALLOW ECF</b> for a correct calculation to 2 DP if:</p> <ul style="list-style-type: none"> <li>%s have been used with wrong isotopes i.e. <math>\frac{(63 \times 30.83) + (65 \times 69.17)}{100} \rightarrow \mathbf{64.38}</math></li> </ul> <p><b>OR</b></p> <ul style="list-style-type: none"> <li>decimal places for <b>ONE</b> % have been transposed, i.e. 69.71 <math>\rightarrow</math> <b>63.96</b>; 30.38 <math>\rightarrow</math> <b>63.32</b></li> </ul>

Question		Answer	Marks	AO element	Guidance
(b)	(ii)	<p><b>FIRST CHECK ANSWER ON THE ANSWER LINE</b>  <b>If answer = <math>3.97 \times 10^{22}</math> (from 63.62) award 2 marks</b>  <b>If answer = <math>3.98 \times 10^{22}</math> (from 63.5) award 2 marks</b></p> <p>-----</p> <p><b>Using 63.62:</b> correct <math>A_r</math> of Cu from 21(b)(i)  See bottom of answer zone</p> $n(\text{Cu}) = \frac{5.00 \times 0.840}{63.62} = \frac{4.2}{63.62} = 0.066(0) \text{ (mol)} \checkmark$ <p>Cu atoms = <math>0.0660 \times 6.02 \times 10^{23} = \mathbf{3.97 \times 10^{22}}</math> <math>\checkmark</math>  <i>Must be calculated in standard form AND to 3 SF</i></p> <p><b>OR</b>-----</p> <p><b>Using 63.5:</b> <math>A_r</math> of Cu from periodic table</p> $n(\text{Cu}) = \frac{5.00 \times 0.840}{63.5} = \frac{4.2}{63.5} = 0.0661 \text{ (mol)} \checkmark$ <p>Cu atoms = <math>0.0661 \times 6.02 \times 10^{23} = \mathbf{3.98 \times 10^{22}}</math> <math>\checkmark</math>  <i>Must be calculated in standard form AND to 3 SF</i></p>	2	AO2.2 x2	<p>If there is an alternative answer, check to see if there is any <b>ECF</b> credit possible  <b>SEE</b> answer from <b>21b(i)</b> at bottom of answer zone</p> <p><b>ALLOW</b> correct answer from 3 SF up to calculator value of 0.06601697579</p> <p><b>ALLOW</b> incorrect <math>n(\text{Cu}) \times 6.02 \times 10^{23}</math> correctly calculated to 3 SF <b>AND</b> in standard form  For <b>ECF</b>, <math>A_r</math> <b>must</b> have been used for <math>n(\text{Cu})</math></p> <p>-----</p> <p><b>ALLOW</b> correct answer from 3 SF up to calculator value of 0.06614173228</p> <p><b>ALLOW</b> incorrect <math>n(\text{Cu}) \times 6.02 \times 10^{23}</math> correctly calculated to 3 SF <b>AND</b> in standard form  For <b>ECF</b>, <math>A_r</math> <b>must</b> have been used for <math>n(\text{Cu})</math></p> <p>-----</p> <p><b>Common errors</b>  <b>Using 63.62:</b>  <math>3.984 \times 10^{22}</math> 1 mark (SF)  <math>4.73 \times 10^{22}</math> 1 mark (ECF: omitting 0.840)  <b>Using 63.5:</b>  <math>3.982 \times 10^{22}</math> 1 mark (SF)  <math>4.74 \times 10^{22}</math> 1 mark (ECF: omitting 0.840)</p>
(c)	(i)	$\text{NiO} + 2\text{HNO}_3 \rightarrow \text{Ni}(\text{NO}_3)_2 + \text{H}_2\text{O} \checkmark$	1	AO1.2	<p><b>ALLOW</b> multiples</p> <p><b>IGNORE</b> state symbols (even if wrong)</p>





Question		Answer	Marks	AO element	Guidance	
22	(a)	<p><b>Initial ratios</b> Cr, <math>\frac{19.51}{52.0}</math> ; Cl, <math>\frac{39.96}{35.5}</math> ; H, <math>\frac{4.51}{1.0}</math> ; O, <math>\frac{36.02}{16.0}</math></p> <p><b>OR</b> Cr, 0.375; Cl, 1.126; H, 4.51; O, 2.25 ✓</p> <p><b>Whole number ratios</b> Cr, 1; Cl, 3; H, 12; O, 6 ✓</p> <p><b>Formula with water of crystallisation</b> CrCl<sub>3</sub>•6H<sub>2</sub>O ✓</p>	3	AO1.2  AO1.2  AO2.2	<p><b>NOTE:</b> If only the correct answer of CrCl<sub>3</sub>•6H<sub>2</sub>O is seen with no working, award 1 mark only</p> <p><b>IF</b> there is no whole number ratio, <b>ALLOW</b> empirical formula: CrCl<sub>3</sub>H<sub>12</sub>O<sub>6</sub></p> <p><b>ALLOW ECF</b> from incorrect whole number ratio, provided <b>ONLY Cl incorrect AND</b> 6H<sub>2</sub>O, e.g. CrCl<sub>2</sub>•6H<sub>2</sub>O</p>	
	(b)	(i)	$\frac{2 \times 0.005}{0.58} \times 100 = 1.72\% \checkmark$	1	AO2.8	<b>ALLOW</b> 2% <b>OR</b> 1.7% up to calculator value of 1.724137931
	(b)	(ii)	Use balance weighing to 3/more decimal places  <b>OR</b>  Use a larger mass/amount <input type="checkbox"/> ✓	1	AO3.3	<p><b>ALLOW</b> more precise/more accurate/ more sensitive/higher resolution/smaller division</p> <p><b>IGNORE</b> 'less error/smaller interval balance'</p> <p><b>IGNORE</b> any reference to lid on crucible (<i>water can't escape</i>)</p> <p><b>IGNORE</b> 'weigh straight after heating'</p> <p><b>IGNORE</b> idea of repeating the experiment/ taking an average/ getting concordant results /larger sample size, etc.</p>

Question		Answer	Marks	AO element	Guidance
(b)	(iii)	Heat to constant mass ✓	1	AO3.4	<b>ALLOW</b> response that implies heating to constant mass, e.g. Heat again until the mass does not change  <b>IGNORE</b> 'heat for longer' <i>Needs link to constant mass</i>
(c)		<p><b>FIRST CHECK ANSWER ON THE ANSWER LINE</b>  <b>If answer = 24.8 (cm<sup>3</sup>) award 3 marks</b></p> <p>-----</p> $n(\text{NaOH}) = 0.124 \times \frac{25.0}{1000} = 3.1(0) \times 10^{-3} \text{ (mol) } \checkmark$ $n(\text{H}_2\text{SO}_4) = \frac{3.10 \times 10^{-3}}{2} = 1.55 \times 10^{-3} \text{ (mol) } \checkmark$ $V(\text{H}_2\text{SO}_4) = 1.55 \times 10^{-3} \times \frac{1000}{6.25 \times 10^{-2}} = 24.8 \text{ (cm}^3\text{) } \checkmark$	3	AO2.8 x3	<p><b>ALLOW ECF</b> from <math>\frac{n(\text{NaOH})}{2}</math></p> <p><b>ALLOW ECF</b> from <math>n(\text{H}_2\text{SO}_4) \times \frac{1000}{6.25 \times 10^{-2}}</math></p>
(d)		<p>Element oxidised: aluminium/Al 0 to +3 ✓</p> <p>Element reduced: hydrogen/H/H<sup>+</sup> +1 to 0 ✓</p>	2	AO1.1 AO1.2	<p><b>MAX 1 mark</b> if no '+' sign for oxidation number</p> <p><b>ALLOW</b> 3+</p> <p><b>ALLOW</b> 1+</p> <p><b>ALLOW</b> H<sub>2</sub> for hydrogen</p> <p><b>ALLOW</b> 1 mark for all oxidation numbers correct, but oxidised and reduced the wrong way around</p> <p><b>IGNORE</b> numbers around equation <i>i.e. treat as rough working</i></p>
		<b>Total</b>	<b>14</b>		



Question		Answer	Marks	AO element	Guidance	
	(c)	(i)	Silver nitrate <b>OR</b> AgNO <sub>3</sub> ✓	1	AO1.1	<b>ALLOW</b> Ag <sup>+</sup> <b>IF</b> name correct, <b>IGNORE</b> an incorrect formula  <b>IGNORE</b> acidified/HNO <sub>3</sub>
	(c)	(ii)	Chloride: white (precipitate) <b>AND</b> Bromide: cream (precipitate) <b>AND</b> iodide: yellow (precipitate) ✓	1	AO1.1	All <b>three</b> required for the mark
			<b>Total</b>	<b>6</b>		

Question		Answer	Marks	AO element	Guidance
24	(a)	<p><b>FIRST, CHECK THE ANSWER ON ANSWER LINE</b>  <b>IF <math>\Delta_r H = -58.5</math> (kJ mol<sup>-1</sup>) award 4 marks</b></p> <p>-----</p> <p><b>Energy released in J OR kJ</b></p> <p><math>= 100.0 \times 4.18 \times 10.5 = 4389</math> (J) <b>OR</b> 4.389 (kJ) ✓</p> <p><b>Correctly calculates <math>n(\text{Pb}(\text{NO}_3)_2)</math></b></p> <p><math>= 1.50 \times \frac{50}{1000} = 0.075(0)</math> (mol) ✓</p> <p><b><math>\Delta H</math> value in J OR kJ</b>  <b>Answer <i>MUST</i> divide energy by <math>n(\text{Pb}(\text{NO}_3)_2)</math></b></p> <p><math>(-)\frac{4389}{0.0750}</math> <b>OR</b> <math>(-)</math>58520 (J)</p> <p><b>OR</b></p> <p><math>(-)\frac{4.389}{0.0750}</math> <b>OR</b> <math>(-)</math>58.52 (kJ) ✓</p> <p><i>(Sign ignored and/or more than 3 SF)</i></p> <p>Correct <math>\Delta_r H</math> in kJ <b>AND</b> – sign <b>AND</b> 3 SF  <math>= -58.5</math> (kJ mol<sup>-1</sup>) ✓</p>	4		<p><b>FULL ANNOTATIONS MUST BE USED</b></p> <p>-----</p> <p>AO2.4 <b>ALLOW</b> 4390 J; 4.39 kJ  <b>DO NOT ALLOW</b> less than 3 SF  <b>IGNORE</b> units  <i>i.e. <b>ALLOW</b> correctly calculated number in J <b>OR</b> kJ</i></p> <p>AO2.4</p> <p><b>ALLOW ECF</b> from <math>n(\text{Pb}(\text{NO}_3)_2)</math> <b>AND/OR</b> Energy  <b>ALLOW</b> 58500 (from 4390)</p> <p>AO2.8 <b>IGNORE</b> absence of – sign and 3 SF requirement</p> <p>AO2.8 Final mark requires – sign, kJ <b>AND</b> 3 SF  <b>Note:</b> From 4390 J, <math>\Delta_r H = -58.5</math> (kJ mol<sup>-1</sup>) (SAME)</p> <p>-----</p> <p><b>Common error</b>  –29.3 3 marks (50 g instead of 100 g in <math>mc\Delta T</math>)</p>

Question		Answer	Marks	AO element	Guidance
	(b)	$\text{Pb}^{2+}(\text{aq}) + 2\text{I}^{-}(\text{aq}) \rightarrow \text{PbI}_2(\text{s}) \checkmark$ <p><b>State symbols required</b></p>	1	AO2.7	<p><b>ALLOW</b> <math>\text{Pb}^{+2}(\text{aq})</math></p> <p><b>IGNORE</b> spectator ions, <math>\text{K}^{+}(\text{aq})</math> and <math>2\text{NO}_3^{-}(\text{aq})</math> on both sides</p>
	(c)	<p><b>FIRST, CHECK ANSWER ON ANSWER LINE IF [KI(aq)] rounds to 3.3 mol dm<sup>-3</sup></b> e.g. 3.30, 3.33, 3.3 recurring</p> <p>-----</p> <p><b>Method 1</b> [KI(aq)] for complete reaction  <math display="block">= 2 \times 0.0750 = 0.150 \text{ mol} \times \frac{1000}{50} = 3 \text{ (mol dm}^{-3}\text{)} \checkmark</math>  10% greater gives <math>3 \times 1.1 = 3.3(0) \checkmark</math></p> <p><b>OR</b>-----</p> <p><b>Method 2</b> <math>n(\text{KI}(\text{aq}))</math> required = <math>2.2 \times 0.0750 = 0.165 \text{ mol} \checkmark</math>  <math display="block">[\text{KI}(\text{aq})] = 0.165 \times \frac{1000}{50} = 3.3(0) \text{ (mol dm}^{-3}\text{)} \checkmark</math></p>	2	AO2.8 x2	<p><b>ALLOW ECF</b> from incorrect <math>n(\text{Pb}(\text{NO}_3)_2)</math> from <b>24(a)</b> <b>BUT</b> if <b>(a)</b> is incorrect but 0.0750 used here, treat as a fresh start and <b>IGNORE</b> response from <b>24(a)</b></p> <p><b>ALLOW</b> 2 marks for 3.3/3.3 recurring <i>Attempt at increasing concentration by 10%</i>  <math display="block">= 2 \times 0.0750 = 0.150 \text{ mol} \times \frac{1000}{45} = 3.33 \text{ (mol dm}^{-3}\text{)}</math></p> <p><b>ALLOW ECF</b> from incorrect <math>n(\text{KI})</math></p> <p>-----</p> <p><b>Common errors</b></p> <p>3            1 mark (Correct for KI with no extra 10%)  1.65        1 mark (no factor of 2 used for KI)  2.7         1 mark (10% less rather than 10% more)  2.73/2.72 1 mark (10% increase in volume: 55 cm<sup>3</sup>)</p>
		<b>Total</b>	<b>7</b>		

Question		Answer	Marks	AO element	Guidance
25	(a)	<p><b>EQUILIBRIUM CONDITIONS</b> <span style="float: right;"><b>3 MAX</b></span>  <b>4 marking points</b> → 3 max ✓✓✓  <i>Mark first three <b>CORRECT</b> responses seen</i></p> <p><b>Temperature:</b>  (Forward) reaction is exothermic/<math>\Delta H</math> is negative  <b>OR</b> (Forward) reaction gives out heat ✓</p> <p><b>Pressure:</b>  Right-hand side has fewer (gaseous) moles  <b>OR</b> 3 (gaseous) moles form 2 (gaseous) moles ✓</p> <p><b>Equilibrium shift</b>  Correct equilibrium shift in terms of <b>temperature</b> ✓  Correct equilibrium shift in terms of <b>pressure</b> ✓</p> <p>-----</p> <p><b>INDUSTRIAL CONDITIONS</b>  Low temperature gives a slow rate/slower reaction  <b>OR</b> high temperatures needed to increase rate ✓□</p> <p>(High) pressure provides a safety risk  <b>OR</b>  (High) pressure is expensive (to generate)  /uses a lot of energy ✓□</p>	5	<p>AO3.1 x2</p> <p>AO3.2 x1</p> <p>-----</p> <p>AO1.2 x2</p>	<p><b>FULL ANNOTATIONS MUST BE USED</b>  -----</p> <p><b>ALLOW</b> suitable alternatives for 'towards right',  e.g.: towards <math>\text{SO}_3</math>/products  <b>OR</b> in forward direction <b>OR</b> 'favours the right'</p> <p><b>ALLOW reverse</b> reaction is endothermic  /<math>\Delta H</math> is positive/takes in heat</p> <p>For moles, <b>ALLOW</b> molecules/particles</p> <p><b>ORA</b> for reverse reaction</p> <p><b>IGNORE</b> responses in terms of activation energy</p> <p><b>ALLOW</b> high pressure is dangerous/explosive</p> <p><b>ALLOW</b> 'These conditions are expensive'  Statement subsumes <b>pressure</b> as 'these' will apply  to <b>pressure</b> (required for this mark) and temperature</p> <p><b>ALLOW ORA</b>  e.g. Lower pressure → less danger/uses less energy</p> <p><b>IGNORE</b> 'It's expensive'  Link with pressure required</p>

Question		Answer	Marks	AO element	Guidance
	(b)	<p><b>Value of <math>K_c</math></b>                      <b>1 mark</b>  <math>K_c</math> is small <b>OR</b> <math>K_c &lt; 1</math>  <b>AND</b> equilibrium (position) is towards left ✓</p>	4		<p><b>FULL ANNOTATIONS MUST BE USED</b>            -----  <b>ALLOW</b> suitable alternatives for ‘towards left,            e.g.: towards <math>\text{SO}_2/\text{O}_2</math> <b>OR</b> towards reactants  <b>OR</b> in reverse direction <b>OR</b> ‘favours the left</p>
		<p><b>Calculation: FIRST CHECK ANSWER</b>  <b>IF <math>[\text{SO}_3] = 0.876</math> OR <math>0.88</math> (mol dm<sup>-3</sup>)</b>  <b>award all 3 marks available for calculation</b>            -----</p> <p><b><math>K_c</math> expression</b>                                      <b>1 mark</b>  <math display="block">\frac{[\text{SO}_3]^2}{[\text{SO}_2]^2[\text{O}_2]} \text{ OR } \frac{[\text{SO}_3]^2}{2.00^2 \times 1.20} \checkmark</math></p> <p><b>Evaluation of <math>K_c</math> <math>[\text{SO}_2]^2[\text{O}_2]</math></b>                      <b>1 mark</b>  <math display="block">K_c [\text{SO}_2]^2[\text{O}_2] = 0.160 \times 2.00^2 \times 1.20</math> <math display="block">= 0.768 \checkmark</math></p> <p><b>Calculation of <math>[\text{SO}_3]</math></b>  <b>ONLY available from correct evaluation for 2nd mark</b>  <math display="block">[\text{SO}_3] = \sqrt{0.160 \times 2.00^2 \times 1.20}</math> <math display="block">= 0.876 \text{ (mol dm}^{-3}\text{)} \checkmark</math></p>		<p>AO1.2                      Square brackets required in <math>K_c</math> expression  <b>ALLOW ECF</b> from <math>\frac{[\text{SO}_3]}{[\text{SO}_2]^2[\text{O}_2]}</math>, i.e. no <math>[\text{SO}_3]^2</math></p> <p>AO2.6                      <b>ALLOW 0.77 (2 SF)</b></p> <p>AO2.6                      <b>ALLOW 0.88 (2 SF)</b> up to calculator value of 0.876356092 correctly rounded</p> <p>AO2.6                      <b>IF <math>K_c</math> expression is inverted 2nd and 3rd marks are available by ECF:</b>  <math display="block">[\text{SO}_3]^2 = \frac{2.00^2 \times 1.20}{0.160} \text{ OR } 30 \checkmark</math> <math display="block">[\text{SO}_3] = \sqrt{30} = 5.48 \text{ OR } 5.5 \checkmark</math></p> <p>Any other <math>K_c</math> expression → <b>NO MARKS</b>,            e.g. <math>\frac{[\text{SO}_3]^2}{[\text{SO}_2]^2 + [\text{O}_2]} \rightarrow \sqrt{0.832} \rightarrow 0.912</math>      <b>NO marks</b></p>	
		<b>Total</b>	<b>9</b>		





Question		Answer	Marks	AO element	Guidance
(c)	(ii)	Compound E: $\begin{array}{c} \text{H} \quad \text{CH}_3 \\   \quad   \\ \text{Br}-\text{C}-\text{C}-\text{Br} \\   \quad   \\ \text{H} \quad \text{CH}_3 \end{array} \quad \checkmark$ Stage 1: Compound E: Bromine/Br <sub>2</sub> ✓  Stage 2: NaOH/KOH OR OH <sup>-</sup> ✓ <i>Only award if intermediate contains at least one halogen atom</i>	3		<b>For structures:</b> <b>ALLOW</b> correct structural <b>OR</b> skeletal <b>OR</b> displayed formula <b>OR</b> mixture of the above  <b>ALLOW</b> dichloro/diiodo compound  <b>IGNORE</b> connectivity of bonds to CH <sub>3</sub>  AO3.2  AO3.1 <b>ALLOW</b> chlorine/Cl <sub>2</sub> <b>OR</b> iodine/I <sub>2</sub> <b>IGNORE</b> conditions, e.g. u.v.  AO3.1 <b>DO NOT ALLOW</b> H <sub>2</sub> O <b>IGNORE</b> conditions  <b>NOTE:</b> Max of <b>2 marks</b> available for <b>monobrominated</b> intermediate  <b>1 mark</b> Reagent: HBr <b>AND</b> Intermediate: CH <sub>3</sub> C(CH <sub>3</sub> ) <sub>2</sub> Br <b>OR</b> BrCH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>  <b>1 mark</b> Intermediate: CH <sub>3</sub> C(CH <sub>3</sub> ) <sub>2</sub> Br <b>OR</b> BrCH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub> <b>AND</b> Reagent: NaOH
<b>Total</b>			<b>8</b>		

**OCR (Oxford Cambridge and RSA Examinations)**  
**1 Hills Road**  
**Cambridge**  
**CB1 2EU**

**OCR Customer Contact Centre**

**Education and Learning**

Telephone: 01223 553998

Facsimile: 01223 552627

Email: [general.qualifications@ocr.org.uk](mailto:general.qualifications@ocr.org.uk)

[www.ocr.org.uk](http://www.ocr.org.uk)

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

**Oxford Cambridge and RSA Examinations**  
**is a Company Limited by Guarantee**  
**Registered in England**  
**Registered Office; 1 Hills Road, Cambridge, CB1 2EU**  
**Registered Company Number: 3484466**  
**OCR is an exempt Charity**

**OCR (Oxford Cambridge and RSA Examinations)**  
**Head office**  
**Telephone: 01223 552552**  
**Facsimile: 01223 552553**

© OCR 2016

