

A-level Chemistry


Paper 2 (7405/2): Organic and Physical Chemistry
Mark scheme

7405
Specimen paper


Version 0.5

Question	Marking guidance	Mark	AO	Comments
01.1	Consider experiments 1 and 2: [B constant] [A] increases $\times 3$: rate increases by 3^2 therefore 2nd order with respect to A	1	AO3 1a	
	Consider experiments 2 and 3: [A] increases $\times 2$: rate should increase $\times 2^2$ but only increases $\times 2$ Therefore, halving [B] halves rate and so 1st order with respect to B	1	AO3 1a	
	Rate equation: $\text{rate} = k[\text{A}]^2[\text{B}]$	1	AO3 1b	
01.2	$\text{rate} = k[\text{C}]^2[\text{D}]$ therefore $k = \text{rate} / [\text{C}]^2[\text{D}]$	1	AO2h	
	$k = \frac{7.2 \times 10^{-4}}{(1.9 \times 10^{-2})^2 \times (3.5 \times 10^{-2})} = 57.0$	1	AO2h	Allow consequential marking on incorrect transcription
	$\text{mol}^{-2} \text{dm}^6 \text{s}^{-1}$	1	AO2h	Any order
01.3	$\text{rate} = 57.0 \times (3.6 \times 10^{-2})^2 \times 5.4 \times 10^{-2} = 3.99 \times 10^{-3} \text{ (mol dm}^{-3} \text{ s}^{-1})$ OR Their $k \times (3.6 \times 10^{-2})^2 \times 5.4 \times 10^{-2}$	1	AO2h	

01.4	Reaction occurs when molecules have $E \geq E_a$	1	AO1a	
	Doubling T causes <u>many</u> more molecules to have this E	1	AO1a	
	Whereas doubling [E] only doubles the number with this E	1	AO1a	
01.5	$E_a = RT(\ln A - \ln k)/1000$	1	AO1b	Mark is for rearrangement of equation and factor of 1000 used correctly to convert J into kJ
	$E_a = 8.31 \times 300 (23.97 - (-5.03))/1000 = 72.3 \text{ (kJ mol}^{-1}\text{)}$	1	AO1b	

Question	Marking guidance	Mark	AO	Comments
02.1	<p data-bbox="248 448 577 480">Gradient drawn on graph</p> 	1	AO3 1a	Line must touch the curve at 0.012 but must not cross the curve.




02.2	<p>Stage 1: Rate of reaction when concentration = $0.0120 \text{ mol dm}^{-3}$ From the tangent Change in [butadiene] = $-0.0160 - 0$ and change in time = $7800 - 0$ Gradient = $-(0.0160 - 0)/(7800 - 0) = -2.05 \times 10^{-6}$ Rate = $2.05 \times 10^{-6} \text{ (mol dm}^{-3} \text{ s}^{-1})$</p> <p>Stage 2: Comparison of rates and concentrations Initial rate/rate at $0.0120 = (4.57 \times 10^{-6})/(2.05 \times 10^{-6}) = 2.23$ Initial concentration/concentration at point where tangent drawn = $0.018/0.012 = 1.5$</p> <p>Stage 3: Deduction of order If order is 2, rate should increase by factor of $(1.5)^2 = 2.25$ this is approximately equal to 2.23 therefore order is 2nd with respect to butadiene</p>			<p>Extended response</p> <p>1 AO3 1a</p> <p>1 AO3 1a</p> <p>1 AO3 1a</p> <p>1 AO3 1a</p> <p>1 AO3 1b</p> <p>Marking points in stage 2 can be in either order</p>
------	--	--	--	--

Question	Marking guidance	Mark	AO	Comments
03.1	2,2,4-trimethylpentane	1	AO1a	
03.2	5	1	AO2b	
03.3	$C_{20}H_{42} \longrightarrow C_8H_{18} + 2C_3H_6 + 3C_2H_4$	1	AO2b	
03.4	Mainly alkenes formed	1	AO1b	
03.5	4 (monochloro isomers) $ \begin{array}{ccccccc} & & CH_3 & H & & CH_3 & \\ & & & & & & \\ H_3C & - & C & - & C & - & C & - & CH_3 \\ & & & & & & & & \\ & & H & & Cl & & CH_3 & & \end{array} $	1	AO2a	
03.6		1	AO2a	

Question	Marking guidance	Mark	AO	Comments
04.1	3-methylbutan-2-ol	1	AO1a	
04.2	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\text{C}-\text{C}-\text{CH}_3 \\ \quad \\ \text{H} \quad \text{O} \end{array} $	1	AO2g	Allow $(\text{CH}_3)_2\text{CHCOCH}_3$
04.3	Elimination	1	AO1a	
04.4	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\text{C}=\text{C}-\text{CH}_3 \\ \quad \quad \\ \quad \quad \text{H} \end{array} $	1	AO2g	Allow $(\text{CH}_3)_2\text{C}=\text{CHCH}_3$
	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\text{C}-\text{C}=\text{CH}_2 \\ \quad \\ \text{H} \quad \text{H} \end{array} $	1	AO2g	Allow $(\text{CH}_3)_2\text{CHCH}=\text{CH}_2$

04.5	Position	1	AO1a	
04.6	C B A	1	AO3 1b	
04.7	$\begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\text{C}-\text{CH}_2\text{CH}_3 \\ \\ \text{OH} \end{array}$	1	AO2g	Allow $(\text{CH}_3)_2\text{C}(\text{OH})\text{CH}_2\text{CH}_3$
04.8	$\begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\text{C}-\text{CH}_2\text{OH} \\ \\ \text{CH}_3 \end{array}$	1	AO2e	Allow $(\text{CH}_3)_3\text{CCH}_2\text{OH}$

Question	Marking guidance	Mark	AO	Comments
05.1	Secondary	1	AO1a	
05.2	Nitrogen and oxygen are very electronegative Therefore, C=O and N–H are polar Which results in the formation of a hydrogen bond between O and H In which a lone pair of electrons on an oxygen atom is strongly attracted to the δ^+ H	1 1 1 1	AO1a AO1a AO1a AO1a	

Question	Marking guidance	Mark	AO	Comments
06.1	 <p style="text-align: center;">glycine threonine lysine</p>	1	AO2a	
06.2		1	AO2a	
06.3		1	AO2a	Allow $(\text{CH}_3)_3\text{N}^+\text{CH}_2\text{COOH} \quad (\text{Br}^-)$
06.4	2-amino-3-hydroxybutanoic acid	1	AO2a	


06.5	$\begin{array}{c} + \\ \text{NH}_3 \\ \\ (\text{CH}_2)_4 \\ \\ \text{H}_3\text{N}^+ - \text{C} - \text{COOH} \\ \\ \text{H} \end{array}$	1	AO2a	
------	--	---	------	--

Question	Marking guidance	Mark	AO	Comments
07.1	$\begin{array}{c} \text{H} \quad \text{CH}_3 \\ \quad \\ \text{C} = \text{C} \\ \quad \\ \text{CH}_3 \quad \text{Cl} \end{array}$ <p>Addition</p>	1 1	AO1a AO1a	
07.2	$\begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{HO}-\text{C}-\text{C}-\text{OH} \\ \quad \\ \text{CH}_3 \quad \text{CH}_3 \end{array}$ $\begin{array}{c} \text{O} \quad \text{CH}_3 \quad \text{H} \quad \text{O} \\ \quad \quad \quad \\ \text{HO}-\text{C}-\text{C}-\text{C}-\text{C}-\text{OH} \\ \quad \quad \\ \text{CH}_3 \quad \text{H} \end{array} \quad \text{OR} \quad \begin{array}{c} \text{O} \quad \text{CH}_3 \quad \text{H} \quad \text{O} \\ \quad \quad \quad \\ \text{Cl}-\text{C}-\text{C}-\text{C}-\text{C}-\text{Cl} \\ \quad \quad \\ \text{CH}_3 \quad \text{H} \end{array}$	1 1	AO2e AO2e	
07.3	<p>Q is biodegradable</p> <p>Polar C=O group or δ^+ C in Q (but not in P)</p> <p>Therefore, can be attacked by nucleophiles (leading to breakdown)</p>	1 1 1	AO2g AO2c AO2c	


Question	Marking guidance	Mark	AO	Comments
08.1	2-deoxyribose	1	AO1a	
08.2	Base A Top N–H forms hydrogen bonds to lone pair on O of guanine The lone pair of electrons on N bonds to H–N of guanine A lone pair of electrons on O bonds to lower H–N of guanine	1 1 1 1	AO3 1b AO2a AO2a AO2a	If Base B stated, allow 1 mark only for response including hydrogen bonding Allow all 4 marks for a correct diagram showing the hydrogen bonding Students could also answer this question using labels on the diagram
08.3	Allow either of the nitrogen atoms with a lone pair NOT involved in bonding to cytosine	1	AO2a	
08.4	Use in very small amounts / target the application to the tumour	1	AO2e	

Question	Marking guidance	Mark	AO	Comments
09.1	<p>(nucleophilic) addition-elimination</p> <p>M2</p> <p>M1</p> <p>M3</p> <p>M4 for 3 arrows and lp</p>	1	AO1a	<p>Not electrophilic addition-elimination</p> <p>Allow C₆H₅ or benzene ring</p> <p>Allow attack by :NH₂C₆H₅</p> <p>M2 not allowed independent of M1, but allow M1 for correct attack on C⁺</p> <p>M3 for correct structure <u>with charges</u> but lone pair on O is part of M4</p> <p>M4 (for three arrows and lone pair) can be shown in more than one structure</p>

09.2	The minimum quantity of hot water was used: To ensure the hot solution would be saturated / crystals would form on cooling	1	AO1b	Allow better drying but not water squeezed out
	The flask was left to cool before crystals were filtered off: Yield lower if warm / solubility higher if warm	1	AO1b	
	The crystals were compressed in the funnel: Air passes through the sample not just round it	1	AO1b	
	A little cold water was poured through the crystals: To wash away soluble impurities	1	AO1b	
09.3	Water	1	AO3 1b	Do not allow unreacted reagents
	Press the sample of crystals between filter papers	1	AO3 2b	Allow give the sample time to dry in air
09.4	M_r product = 135.0	1	AO2h	Answer must be given to this precision
	Expected mass = $5.05 \times \frac{135.0}{93.0} = 7.33$ g	1	AO2h	
	Percentage yield = $\frac{4.82}{7.33} \times 100 = 65.75 = 65.8(\%)$	1	AO1b	

09.5	 <p>OR</p> $\text{C}_6\text{H}_5\text{NHCOCH}_3 + \text{NO}_2^+ \rightarrow \text{C}_6\text{H}_4(\text{NHCOCH}_3)\text{NO}_2 + \text{H}^+$	1	AO2c	
09.6	Electrophilic substitution	1	AO1a	
09.7	Hydrolysis	1	AO3 1a	
09.8	Sn/HCl	1	AO1b	Ignore acid concentration; allow Fe/HCl

Question	Marking guidance	Mark	AO	Comments								
11.1	$\text{CH}_3\text{CH}_2\text{COCH}_3 + 2[\text{H}] \longrightarrow \text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$	1	AO1b									
11.2	<p>This question is marked using levels of response. Refer to the Mark Scheme Instructions for Examiners for guidance on how to mark this question.</p> <table border="1" data-bbox="235 646 1234 1423"> <tbody> <tr> <td data-bbox="235 646 405 821">Level 3 5–6 marks</td> <td data-bbox="405 646 1234 821"> <p>All stages are covered and the explanation of each stage is generally correct and virtually complete.</p> <p>Answer is communicated coherently and shows a logical progression from stage 1 to stage 2 then stage 3.</p> </td> </tr> <tr> <td data-bbox="235 821 405 1082">Level 2 3–4 marks</td> <td data-bbox="405 821 1234 1082"> <p>All stages are covered but the explanation of each stage may be incomplete or may contain inaccuracies OR two stages are covered and the explanations are generally correct and virtually complete.</p> <p>Answer is mainly coherent and shows progression from stage 1 to stage 3.</p> </td> </tr> <tr> <td data-bbox="235 1082 405 1329">Level 1 1–2 marks</td> <td data-bbox="405 1082 1234 1329"> <p>Two stages are covered but the explanation of each stage may be incomplete or may contain inaccuracies, OR only one stage is covered but the explanation is generally correct and virtually complete.</p> <p>Answer includes isolated statements but these are not presented in a logical order or show confused reasoning.</p> </td> </tr> <tr> <td data-bbox="235 1329 405 1423">Level 0 0 marks</td> <td data-bbox="405 1329 1234 1423">Insufficient correct chemistry to gain a mark.</td> </tr> </tbody> </table>	Level 3 5–6 marks	<p>All stages are covered and the explanation of each stage is generally correct and virtually complete.</p> <p>Answer is communicated coherently and shows a logical progression from stage 1 to stage 2 then stage 3.</p>	Level 2 3–4 marks	<p>All stages are covered but the explanation of each stage may be incomplete or may contain inaccuracies OR two stages are covered and the explanations are generally correct and virtually complete.</p> <p>Answer is mainly coherent and shows progression from stage 1 to stage 3.</p>	Level 1 1–2 marks	<p>Two stages are covered but the explanation of each stage may be incomplete or may contain inaccuracies, OR only one stage is covered but the explanation is generally correct and virtually complete.</p> <p>Answer includes isolated statements but these are not presented in a logical order or show confused reasoning.</p>	Level 0 0 marks	Insufficient correct chemistry to gain a mark.	6	1 AO1a 5 AO2a	<p>Indicative Chemistry content</p> <p>Stage 1: Formation of product</p> <ul style="list-style-type: none"> Nucleophilic attack Planar carbonyl group H^- attacks from either side (stated or drawn) <p>Stage 2: Nature of product</p> <ul style="list-style-type: none"> Product of step 1 shown This exists in two chiral forms (stated or drawn) Equal amounts of each enantiomer/racemic mixture formed <p>Stage 3: Optical activity</p> <ul style="list-style-type: none"> Optical isomers/enantiomers rotate the plane of polarised light equally in opposite directions With a racemic/equal mixture the effects cancel
Level 3 5–6 marks	<p>All stages are covered and the explanation of each stage is generally correct and virtually complete.</p> <p>Answer is communicated coherently and shows a logical progression from stage 1 to stage 2 then stage 3.</p>											
Level 2 3–4 marks	<p>All stages are covered but the explanation of each stage may be incomplete or may contain inaccuracies OR two stages are covered and the explanations are generally correct and virtually complete.</p> <p>Answer is mainly coherent and shows progression from stage 1 to stage 3.</p>											
Level 1 1–2 marks	<p>Two stages are covered but the explanation of each stage may be incomplete or may contain inaccuracies, OR only one stage is covered but the explanation is generally correct and virtually complete.</p> <p>Answer includes isolated statements but these are not presented in a logical order or show confused reasoning.</p>											
Level 0 0 marks	Insufficient correct chemistry to gain a mark.											

Question	Marking guidance	Mark	AO	Comments
12.1	HBr OR HCl OR H ₂ SO ₄	1	AO1b	Allow HI or HY
12.2	Electrophilic addition 	1 4	AO1a AO2a	Allow consequential marking on acid in 12.1 and allow use of HY
12.3	The major product exists as a pair of enantiomers The third isomer is 1-bromobutane (minor product) Because it is obtained via primary carbocation	1 1 1	AO2a AO2a AO2a	